


A Brief History of Triangulations

University College Cork

Wednesday, March 25, 2026

Christopher Bishop, Stony Brook University

slides posted at www.math.sunysb.edu/~bishop/lectures






Egyptian triangles



French triangles



American triangles



Irish triangles

triantáin



Fíor iata é an triantán.
Bíonn trí shlios air.

UILLINEACHA AN TRIANTÁIN



Dronuillinn (90°)
amháin a bhíonn
i dtriantán dronuilleach.



I dtriantán maoluilleach
bíonn uillinn amháin níos
mó ná 90° .



I dtriantán géaruilleach
bíonn na trí uillinn níos
lú ná 90° .

SLEASA AN TRIANTÁIN



I dtriantán comhshleasach
bíonn na trí shlios
ar comfhad.



I dtriantán comhchosach
bíonn dhá shlios
ar comfhad.



I dtriantán corrsheasach
bíonn fad éagsúil
sna trí shlios.

CINEÁLACHA TRIANTÁN



Triantán
comhchosach
dronuilleach



Triantán
corrsheasach
dronuilleach



Triantán
comhchosach
maoluilleach



Triantán
corrsheasach
maoluilleach



Triantán
comhchosach
gearuilleach



Triantán
corrsheasach
gearuilleach

Triangles in Irish




Flying triangles




Triangles in art (Picasso)




Triangles in engineering




Triangles in computer graphics
(The Stanford Bunny)




Triangulation of a polygon




Triangulation of a polygon



Triangulation of a polygon




Triangulation of a polygon




Triangulation of a polygon
using only the vertices of the polygon

There are three types of triangulations:


- Triangulation (no extra vertices)
- Steiner triangulation (extra vertices allowed, edges match)
- Triangular dissections (extra vertices allowed, edges need not match)




We can add extra points, called Steiner points




Using these, we get a Steiner triangulation.



Simplex condition: adjacent triangles have matching edges.
Most applications use this condition.



A dissection of a polygon, edges can mis-match




In a **triangulation**, triangles meet only at vertices or full edges.

In a **dissection**, triangles can meet along partial edges.


dissections

Steiner triangulations

triangulations




Does every polygon have a triangulation? (no extra vertices)



Yes. Can prove by induction on number of sides N .

$N = 3$ is obvious. For $N > 3$, it suffices to find one diagonal.

Diagonal = segment connecting two vertices **inside** the polygon.




A diagonal cuts polygon into two pieces each with fewer sides.
Each piece can be triangulated by the induction hypothesis.

Proof that a diagonal always exists:

Claim: every polygon has an interior angle $< 180^\circ$.

Proof that a diagonal always exists:


Claim: every polygon has an interior angle $< 180^\circ$.



Choose line outside P and not parallel to any side.

Proof that a diagonal always exists:


Claim: every polygon has an interior angle $< 180^\circ$.



Choose line outside P and not parallel to any side.
Slide the line until it first hits P .

Proof that a diagonal always exists:


Claim: every polygon has an interior angle $< 180^\circ$.



Choose line outside P and not parallel to any side.
Slide the line until it first hits P .

Proof that a diagonal always exists:

Claim: every polygon has an interior angle $< 180^\circ$.




Choose line outside P and not parallel to any side.

Slide the line until it first hits P .


Intersection must contain a vertex v with angle $< 180^\circ$.

Proof that a diagonal always exists, Case 1:




Connect vertices v_1, v_2 adjacent to v by a segment S .
If this segment S inside P , we are done.

Proof that a diagonal always exists, Case 2:




Otherwise, segment $[v_1, v_2]$ hits other edges.

Proof that a diagonal always exists, Case 2:




Move S towards v , parallel to itself,
until last time it hits an edge not adjacent to v .

Proof that a diagonal always exists, Case 2:




Move S towards v , parallel to itself,
until last time it hits an edge not adjacent to v .

Proof that a diagonal always exists, Case 2:




Move S towards v , parallel to itself,
until last time it hits an edge not adjacent to v .

Proof that a diagonal always exists, Case 2:




Then S contains a vertex w that can be connected to v . QED



Two polygons cut into same set of triangles must have same area.

Wallace–Bolyai–Gerwien Theorem (early 1800's):


Any two polygons with the same area have dissections into similar sets of sub-polygons (up to rotations and translations).



An equi-dissection of an equilateral triangle and a square.

Wallace–Bolyai–Gerwien Theorem (early 1800's):

Any two polygons with the same area have dissections into similar sets of sub-polygons (up to rotations and translations).



An equi-dissection of an equilateral triangle and a square.


Who proved the WBG theorem?

Not clear. Contradictory information given in:


I. Stewart, *From here to infinity*, The Clarendon Press, 1996.

E.N. Giovannini, David Hilbert and the foundations of the theory of plane area, *Arch. Hist. Exact Sci.*, vol 75, 2021, pages 649–698.

Lemma: If P_1 and P_2 have an equi-dissection, and if P_2 and P_3 have an equi-dissection, then so do P_1 and P_3 . (Equi-dissection is transitive.)




Lemma: If P_1 and P_2 have an equi-dissection, and if P_2 and P_3 have an equi-dissection, then so do P_1 and P_3 . (Equi-dissection is transitive.)



Proof: Intersect the two dissections of P_2 . Smaller pieces can be rearranged into either P_1 or P_3 .

Proof of WBG theorem.


Enough to show a polygon of area A is equivalent to a $1 \times A$ rectangle.



Proof of WBG theorem.


It is enough to show a triangle of area a is equivalent to a $1 \times a$ rectangle.

Triangulate P , convert each triangle to a rectangle and stack them.




Proof of WBG theorem.

Claim 1: every triangle is dissection equivalent to some rectangle:




Proof of WBG theorem.

Claim 2: a $t \times s$ rectangle is equivalent to a $(t/2) \times (2s)$ rectangle.



Proof of WBG theorem.

Claim 2: a $t \times s$ rectangle is equivalent to a $(t/2) \times (2s)$ rectangle.




Claim 3: a rectangle of area a is equivalent to a rectangle with side lengths $s \in [1, 2)$ and $t \in (a/2, a]$.

Keep doubling (or halving) sides until this is true.

Proof of WBG theorem.



Claim 4: a rectangle of area a is equivalent to a $1 \times a$ rectangle.

By previous steps we may assume $t \times s$ with $1 \leq s < 2$.




This completes the proof of the WBG theorem.

What about dimension 3?



3D version of a triangle is a tetrahedron = four faces, all triangles

What about dimension 3?




3D version of a polygon is a polyhedron.

So far we have proven

- Every 2D polygon has a triangulation with no extra vertices.
- The Wallace-Bolyai-Gerwien theorem.


Are these true in 3 dimensions?

- Is every polyhedron a union of tetrahedron using no extra vertices?
- Can two equal volume polyhedra be cut into the same set of tetrahedra?





Some polyhedra can be cut into tetrahedra.
A triangular prism can be cut into three tetrahedra.

3D counterexample due to Erich Schönhardt in 1928.



The Schönhardt polyhedron does not contain any tetrahedra defined by four of its vertices.


3D counterexample due to Erich Schönhardt in 1928.




The Schönhardt polyhedron does not contain any tetrahedra defined by four of its vertices.

Proof:

- Let P be the Schönhardt polyhedron.
- Suppose T is a tetrahedron using 4 of the 6 vertices of P .
- T is inside P iff none of its vertices can “see” another one.
- Every vertex v on top side of P , can see one vertex on bottom side.
- If T has 3 vertices on top, the vertex on bottom sees one of these. $\Rightarrow \Leftarrow$
- If T has 2 vertices on top, only one on bottom is “invisible” $\Rightarrow \Leftarrow$





Is the WBG Theorem true in 3 dimensions?
Can we cut a cube into tetrahedra and re-assemble as a pyramid?




This is “**Hilbert’s third problem**”. Ten problems given at the 1900 International Congress of Mathematicians; 23 later appeared in print.

The Dehn Invariant is a “number” associated to each polyhedron.

Dehn proved that dissection equivalent polyhedra have same value.

But unit cube and regular tetrahedron have different Dehn invariants.



Max Dehn



In 1965 Jean-Pierre Sydler proved converse:


Theorem: Two polyhedra are dissection equivalent iff equal same Dehn invariants.

Easier algebraic proof given in 1990 by Johan Dupont and Han Sah.

Proof is related to topology and the homology the 3-D rotation group.

Defining the Dehn invariant:

- “Angles” means dihedral angles between faces. One per edge.



Defining the Dehn invariant:

- The reals form an (infinite dimensional) vector space over the rationals.


Vector space = set with addition and multiplication by some scalar field.

For example:

$$\log 120 = 3 \log 2 + \log 3 + \log 5.$$

Exercise: $\{\log p\}$ are independent over rationals for p prime.

Research problem: is e a rational multiple of π ?



Defining the Dehn invariant:

Given a polyhedron:

- (1) Find a basis for the dihedral angles over the rationals. Add π .
- (2) Write each dihedral angle as a sum of basis elements, i.e., a vector.
- (3) Multiply each vector by corresponding edge length.
- (4) Dehn Invariant = sum vectors over all edges and drop the π coordinate.


Platonic solids have only one angle, so Dehn invariant is 1-dimensional.

Enough to check if dihedral angles are rational multiples of each other.


Dihedral angles: Cube = $\pi/2$, Regular Tetrahedron = $\arccos(1/3)$.

Fact: $\arccos(1/3)$ is not a rational multiple of π .

\Rightarrow Cube and Tetrahedron are not dissection equivalent.



\neq




$$\frac{\arccos(1/3)}{\pi} =$$

0. 391826552030607270170855559222430911316628649439400
287287938298022001408677442246870211826175774513079
526782730495457978237526235003385455266599415972920
905797050905025019933356981299552610344832589848763
539096857387170086489247019208018189353116296066293
190417231448867474699006802327685632777984042777431
659577084905067798797920906188214217781178972740561
610916275778662094588988620484321630438929018193900
355296601454579820496938390427736624997412806048800
79574307427701871242437713360951036300515 ...

Doesn't look rational, but proof given at end of slides.

Surprising historical note:

Hilbert's problem had been solved even before he stated it.



Władysław Kretkowski



Ludwig Birkenmajer

In 1882, when a Polish nobleman Władysław Kretkowski offered a prize of 500 French francs (about \$10,000 today) for its solution.

Awarded by the Polish Academy of Arts and Sciences to a 28-year-old math teacher, Ludwig Birkenmajer, later a professor at Jagiellonian University.

Equidecomposability of polyhedra: a solution of Hilbert's third problem in Kraków before ICM 1900" by D. and K. Ciesielski, *Mathematical Intelligencer* vol 40, 2018.


Side remark:

Equal volume polyhedra can't always be equi-partitioned into tetrahedra.

What if we replace tetrahedra by more general sets?

Can two polyhedra of equal volume be decomposed into finite collections of sets that are the same, up to rotation and translation?

Banach-Tarski paradox (1926): Given any two polyhedra in 3-space, the first can be decomposed into finitely many disjoint subsets, that can be rotated and translated to give the second as their disjoint union.




Banach-Tarski paradox (1926): Given any two polyhedra in 3-space, the first can be decomposed into finitely many disjoint subsets, that can be rotated and translated to give the second as their disjoint union.



Stephen Banach




Alfred Tarski



This violates our intuition that


- (1) volume does not change under translations or rotations,
- (2) the volume of a disjoint union should be the sum of the volumes.



This violates our intuition that

- (1) volume does not change under translations or rotations,
- (2) the volume of a disjoint union should be the sum of the volumes.

We can't define 3D-volume so it has both properties for all sets.




This is why a field called “Measure Theory” exists.


Volume can only be defined for special “measurable sets”.

“Measurable” includes every set you can explicitly describe.

Non-measurable sets require the Axiom of Choice to construct.




For many applications, it is important to have triangulations with “nice” triangles. (“nice” = angles not too close to 0 or 180.)




Good


Bad




equilateral



acute (all < 90)




right (one = 90)




obtuse (some > 90)


non-obtuse




equilateral



acute (all < 90)




right (one = 90)




obtuse (some > 90)

Acute triangles are often better for computer graphics and engineering. For example, in finite elements methods for differential equations, the systems arising from acute triangulations are often easier and faster to solve.



“An essential element of the numerical solution of partial differential equations (PDEs) on general regions is the construction of a grid (mesh) on which to represent the equations in finite form ...it can take orders of magnitude more man-hours to construct the grid than it does to perform and analyze the PDE solution on the grid.”

Quoted from the introduction *Delaunay Mesh Generation*, by Cheng, Dey and Shewchuk, Chapman & Hall, 2013.




The same text points out that motion pictures are now the most economically significant consumers of high quality meshing algorithms.

We want to mesh surfaces using acute triangulations.

Do acute triangulations always exist?

If so, how many acute triangles are needed?

Burago-Zalgaller (1960): all polyhedral surfaces have acute triangulations.



Yuri Burago (1936 -)



Victor Zalgaller (1920 - 2020)

However, their result was a lemma buried inside a Russian language paper on topology and went unnoticed in the West. First reference I know to Burago-Zalgaller in CS papers is in 2004.


They were proving a polyhedral version of the Nash Embedding Theorem.



Nash embedding




John Nash




not John Nash

Computational complexity = number of triangles needed.

- Baker-Grosse-Rafferty (1987) proved every n -gon has a non-obtuse triangulation (weaker than the Burago-Zalgaller theorem).
- Bern-Mitchell-Ruppert (1995): Cn triangles suffice for simple polygons.
- Maehara (2002): any nonobtuse triangulation, can be subdivided to give an acute triangulation \Rightarrow Burago-Zalgaller + linear complexity.





Non-simple polygon = PSLG = Planar Straight Line Graph

- Bishop (2016): $Cn^{2.5}$ triangles suffice for PSLGs.

Some examples need $\simeq n^2$. Sharp power remains unknown.


Defn: A θ -triangulation is one where every angle is $\leq \theta$.

Burago-Zalgaller \Rightarrow every simple polygon has a θ -triangulation for some $\theta < 90^\circ$.

How does θ depends on the polygon?

From here on, “triangulation” means “Steiner triangulation”.

For example, a square has a 72° -triangulation, and this is optimal.



Optimality proven using Euler's formula ($F - E + V = 1$).

F = number of faces, E = number edges, V = number of vertices.

Proof at end of slides.

For general polygons, we have (Bishop, 2026):


Theorem: The optimal angle bound for triangulating an n -gon can be computed in time $O(n)$.

Theorem: The optimal angle bound only depends on the set of angles, not their order or the edge lengths.


Theorem: If P has minimal angle $\geq 36^\circ$ then it has a 72° -triangulation.

Theorem: The optimal angle bound for triangulations is the same as for triangular dissections.


Theorem: An optimal triangulation “usually” exists.




Maximum angle ≈ 74.7482 .




Maximum angle ≈ 70.3590 .




Maximum angle ≈ 67.8690 .



Optimum bound is $= 67.5^\circ$. Can it be attained?




Optimum bound is $= 67.5^\circ$. Can it be attained?
Yes in this case, but not for all polygons.



If a polygon P has an equilateral triangulation, then:

- (1) all angles of P are integer multiples of 60° (P is a 60° -polygon) .
- (2) all triangles are same size.
 - \Rightarrow all edges of P are multiples of this length
 - \Rightarrow length ratios of any two edges is rational





Lemma: every 60° -polygon has a θ -triangulation for every $\theta > 60^\circ$.

But if some edge ratio is irrational no equilateral triangulation exists.

Conclusion: for such polygons no optimal triangulation exists.

Thm: An optimum triangulation exists for every other polygon.

Very, very rough idea behind proofs:



- Given P “approximate” it by a 60° -polygon P' .
- Use conformal map $P' \rightarrow P$ to transfer equilateral triangulation.
- Prove worst angle distortion is near vertices.


Choosing the correct P' is the hard part.

The Future (open problems)

1. Acute Triangulation in Dimension 3:

Question: Does a polyhedron with n vertices have a polynomial sized acute tetrahedralization?


(dihedral angles $< 90^\circ$, number of elements less than a power of n .)



This is pretty hard even for the unit cube.

The “smallest” known acute triangulation of a cube uses **1370** tetrahedra.

Found by VanderZee, Hirani, Zharnitsky and Damrong in 2010, using a computer search. A more conceptual construction with **2715** tetrahedra was given in 2012 by Kopczyński, Pak and Przytycki.



2. Minimal Weight Triangulations


Find triangulation that minimizes total edge length.



$1 \times R$ rectangle with n extra points on right side.

2. Minimal Weight Triangulations

Find triangulation that minimizes total edge length.




$1 \times R$ rectangle with n extra points on right side.

Without Steiner points, all triangulations length $> R \cdot n$.

For general polygon finding (non-Steiner) optimum is NP hard.

2. Minimal Weight Triangulations

Find triangulation that minimizes total edge length.




$1 \times R$ rectangle with n extra points on right side.

With Steiner points, length $\leq R + O(\log n)$ is possible.

What is optimal? Does an optimal triangulation exist?

2. Minimal Weight Triangulations




As x approached d , total length tends to infimum.

A minimal weight Steiner triangulation (MWST) does not exist.

But there are three co-linear points (b, c, d) .

Question: Does a MWST always exist for general polygons?

3. A stronger Wallace–Bolyai–Gerwien Theorem:




The WBG theorem says two equal area polygons can be dissected using the same set of triangles.

Can “dissection” be improved to “triangulation”?

In other words, all triangles meet along full edges.


3. A stronger Wallace–Bolyai–Gerwien Theorem:




Thm (B, 2026): Any two equal area polygons can be triangulated using same set of triangles.


Proof uses ideas from dynamical systems. Boundaries of the polygons play a very important role. What if there is no boundary?



Question: Can two closed polyhedral surfaces with equal area be triangulated using the same set of triangles?



?







Exercise: Prove that $\theta = \arccos(1/3)$ is not a rational multiple of π .

Solution: We give a proof by contradiction. Suppose $\theta = m\pi/n$. Then $\sin(n\theta) = \sin(m\pi) = 0$, and hence $\cos(n\theta) = \pm 1$. Let $z = \cos \theta + i \sin \theta$. Then $z^n = \cos(n\theta) + i \sin(n\theta) = \pm 1$, so z is an algebraic integer (the root of an equation with integer coefficients, $z^n \pm 1$).

Similarly, $(1/z)^n = \pm 1$, so $1/z$ is also an algebraic integer.

Thus $z + 1/z = 2 \cos(\theta) = 2/3$ is an algebraic integer, which is impossible since the only real rational numbers that are algebraic integers are the regular integers. \square

Exercise: square has no triangulation with all angles $< 72^\circ$.

Given a triangulation of square S , define:

- r_k = number of square's corners in k triangles,
- q_k = number of interior triangulation vertices in k triangles,
- s_k = number of non-corner boundary vertices in k triangles.

We have the relations

- $F - E + V = 1$ (Euler's formula),
- $V = \sum_k (q_k + r_k + s_k)$ (every vertex is one of the three types)
- $3F = \sum_k k(q_k + r_k + s_k)$ (triangle corners counted two ways),
- $2E = 3F - \sum_k (r_k + s_k)$ (triangle sides counted two ways.)

Combining these four equations and eliminating variables gives


$$\sum_k (6 - k)q_k + \sum_k (3 - k)r_k + \sum_k (3 - k)s_k = 6,$$

If every angle is $< 72^\circ$ then every interior vertex is on ≥ 6 triangles and every non-corner boundary vertex is on ≥ 3 triangles. Thus

$$2r_1 + r_2 \geq \sum_k (3 - k)r_k \geq 6.$$



Since $r_2 \leq 4$ (the square has four corners), we have $r_1 > 0$. Thus a 90° angle occurs at some corner, a contradiction. \square

Hyperbolic triangulations:



Equilateral triangulation of disk in hyperbolic metric.



Hyperbolic triangulations:



We say a triangulation is equilateral if any two triangles sharing an edge are reflections of each other across that edge.

This works in either geometry (if we know how to define reflections).

Hyperbolic triangulations:




Some compact surfaces have an equilateral triangulation.

Most do not; only finitely many ways to glue together n triangles.

Compact surfaces with equilateral triangles have been intensely studied: related to Grothendieck's theory of *dessins d'enfants*, graph theory, probability, algebraic number theory, Galois theory, ...

Hyperbolic triangulations:





Thm (Bishop-Rempe, 2026): Every non-compact hyperbolic surface has an equilateral triangulation.

Proof combines differential equations and planar acute triangulation.


Many questions about hyperbolic triangulations remain open.

TRIANGULATION FLOWS







A triangle.




Its in-circle.




The central region and three sectors (thin version).




The three sectors are foliated by circular arcs.
Defines flow on a triangulation that stops at boundary or cusp point.




Delaunay triangulation of 10 random points,




The boundary of the triangulation




The central regions.




Propagation lines starting at all cusp points.




Propagation lines identify boundary points; induces tree.
Discontinuous, but piecewise length preserving.




Delaunay triangulation of 30 random points in disk.




The central regions.




Propagation lines starting at all cusp points.




Enlargement 1.




Enlargement 2.




60 points



The central regions.



Propagation lines starting at all cusp points.



Log-log plot of number points created versus n .
Slope ≈ 2.5